

7 - 12 - 6 + 2MgO

| | |
|---|--------|
| Nitrogen (N) | 7,0 % |
| Ammonium N | 4,8 % |
| Organic N | 2,2 % |
| Phosphorus (P ₂ O ₅) | 12,0 % |
| Potassium (K ₂ O) | 6,0 % |
| Calcium (CaO) | 10,0 % |
| Magnesium (MgO) | 2,0 % |
| Sulphur (SO ₃) | 9,0 % |
| Organic Carbon (C) | 21 % |
| Organic Matter | 41 % |
| Aminoacids & Proteins | 15 % |
| Humic & Fulvic acids | 8 % |
| Moisture | 7 % |
| C/N | 3,0 |

| CROPS | RATE | APPLICATION |
|--|--------------------|------------------------------------|
| Fruit tress, Stone fruits, Pome trees, Grapevines | 500-800 kg/ha | From late Summer to late Spring |
| Citrus | 1400-1600 kg/ha | Winter |
| Olives | 500-700 kg/ha | From late Summer to late Spring |
| Vegetables (open field and greenhouse), Greens | 600-1000 kg/ha | Before sowing or transplanting |
| Cereals, Soya, Rice, Beets, Cotton | 400-500 kg/ha | Before sowing |
| Industrial crops | 400-500 kg/ha | Before sowing |
| Ornamentals | 400-500 kg/ha | Before sowing or transplanting |

Batch num.:
Production date:



Storage conditions: It must be stored in well closed containers, in room temperature, at a dry place without moisture, protected from rain, frost and high temperatures.