


Micronutrient mixture in water soluble powder form

# Elem-Mix



EC Fertilizer

Net Weight

- 1 Kg   
5 Kg   
25 Kg

## Contains

Iron	4,0%	(EDTA)
Zinc	4,0%	(EDTA)
Magnesium	2,0%	(Salt)
Manganese	3,0%	(EDTA)
Boron	1,5%	(Salt)
Copper	0,5%	(EDTA)
Molybdenum	0,005%	(Salt)
Sulfur	2,8%	(Salt)

## **APPLICATION RATES**

The following application rates are general. Actual amounts used will depend on the type and the size of plant and on the degree of deficiency.

### **Foliar application**

*Vegetables* : 1 - 4 kg/ha.

*Cereals* : 1 - 2 kg/ha.

*Fruit trees & bushes* : 2 - 5 kg/ha following a program of repeated intensive sprays, (e.g. 4 applications of 500 g dissolved in 300-1,000 l of water/ha).

*Ornamentals* : 1-4 kg/ha.

*Note* : Application rates must never exceed 2 g/ liter of water.

### **Soil application**

*Vegetables* : 4 - 8 kg/ha.

*Cereals* : 4 - 10 kg/ha.

*Fruit trees & bushes* : 4 - 10 kg/ha.

*Ornamentals* : 2 - 10 kg/ha.

Rates should be adjusted according to the age of the plant, variety, soil type and other local conditions.

## **APPLICATIONS - APPLICATION PERIOD**

**Foliar application**: Foliar application is recommended for the quick correction of deficiencies. In case phytotoxic problems (such as scorch) occur in very sensitive crops after foliar application continue with soil application. As to achieve maximum benefits, foliar applications must begin at the beginning of the growing season, as soon as sufficient foliage exists to absorb the spray. In case of sufficient deficiency, applications must be repeated every 7-14 days. When **Elem-lix** is applied foliarly on its own, it is recommended to add a sticking agent in the solution.

**Soil application**: Dissolve in an appropriate amount of water and apply with a course low pressure spray, or apply mixed with liquid fertilizer. Apply to the soil close to plants and tree trunks. Work well into the soil after application, water carefully after each application in order to enhance nutrient uptake.

**COMPATIBILITY**: **Elem-lix** can be mixed with most fertilizers and agrochemicals without causing precipitation or scorch problems. In liquid fertilizers the mixture should be used as soon as possible.

**STORAGE**: Store in its original container tightly closed in a dry place and away from strong light. Keep it out of the reach of children and pets.

**CAUTION**: Do NOT apply to crops suffering stress such as extreme temperatures, strong winds etc.

## **DANGER**



## **Reproductive toxicant (Category 1B)**


### **Hazard statements:**

H360FD: May damage fertility. May damage the unborn child. (Disodium Octaborate Tetrahydrate)

### **Precautionary statements:**

P201: Obtain special instructions before use. P281: Use personal protective equipment as required. P308+P313: IF exposed or concerned: Get medical advice/attention. P405: Store locked up. P501: Dispose of contents/container to ... in accordance with local/regional/national/ international regulation (to be specified).

Produced by



1 Ermou & Theotokopoulou str  
144 52, Metamorphosis, Greece  
www.humofert.gr