Tri-ACTIVE is a growth biostimulant, which ensures the fortified growth of plants and their root system due to the various strains of the fungus *Trichoderma* that it contains. Tri-ACTIVE contains conidia and mycelium from the beneficial strains of Trichoderma, which colonize right after the application of the product the rhizosphere of the plant and promote the growth both of the underground and the overground part of the plants. Furthermore, they increase the assimilation of nutrients and water from the plants and as a result they contribute to a better nutrition of the crops and the enhancement of their resistance against various environmental -stress factors.

BENEFITS

- Accelerates the growth rate of the plants.
- Promotes the growth of a healthy and voluminous root system.
- " Increases the penetration of the root system ensuring the proper establishment of the plants.
- Increases the number of the roots that reach a great depth enhancing greatly the resistance of the crops against drought.
- " Enhances the availability of the nutrients of the soil, such as phosphorus, iron, copper etc.
- " Increases crop vield.
- Fortifies the resistance of the plants against harsh environmental conditions, salinity, infections by pathogens etc.

RECOMMENDED CROPS

Tri-ACTIVE can be applied in vegetables, horticulture, cereals, aromatic plants, legumes, cotton, oilseed rape, sunflower, cardamom, beets, potato, fruit trees, ornamental bushes and flowers.

PRECAUTIONS

During application of the product you must use gloves, mask, safety glasses and protective clothing. In case of contact with skin or clothing, wash thoroughly your skin with soap and water. In case of contact with the eyes, remove any contact lenses you may wear and flush with plenty of water. If irritation persists, seek medical advice. The product must not be used by people who are immunosuppressed.

KEEP OUT OF THE REACH OF CHILDREN

Tri-ACTIVE

Growth Biostimulant

SYNTHESIS

Beneficial microorganisms >1.5x10¹² cfu*/l Manganese (Mn) 0.1 %

*cfu: colony forming units





Net Content: 1 |

Produced by



1 Ermou & Theotokopoulou str., 144 52 Metamorphosis, Greece Tel. +30 210 284 5891, +30 210 285 5061 Fax. +30 210 281 7971

APPLICATION METHOD

Tri-ACTIVE is applied directly in the soil by watering or drenching around the root area. Right after the application it must be incorporated in the soil mechanically (hoeing) or through irrigation of the plants. Tri-ACTIVE is suitable for use through any fertigation system. In addition it can be applied by dipping the seed prior to planting and the plantlet or the cutting before planting-transplanting.

APPLICATION FREQUENCY

It is applied initially at the first irrigation or during sowing, planting or transplanting and then by repeated applications every 4-6 weeks. The application frequency depends on the state of the soil and the crop.

APPLICATION RATE

Soil application: 5 - 10l/ha.

Application in seedlings: Dipping in a solution 5 ml of product in 10 l of water right before planting.

Seed dipping: 10 ml of product per 1 kg of seed. Prepare a solution of appropriate volume in order for the seeds to be soaked thoroughly. Then let the seeds dry before planting them.

Cutting dipping: In a solution 5 ml of product per 10 l of water, dip the cuttings for 10 minutes and then plant them.

COMPATIBILITY

Tri-ACTIVE cannot be mixed with fungicides. It can be mixed with many liquid fertilizers, insecticides, pesticides and bactericides. In any case it is recommended to avoid applying a soil fungicide for at least 7-10 days before or after the application of Tri-ACTIVE.

STORAGE CONDITIONS

The product must be stored in a cool place, without humidity. Store well sealed in its original container. In case you open the product you must use it immediately, otherwise store it well sealed in the refrigerator (4°C). The shelf life of the product in normal room conditions is 8-12 months and 1-2 years at 4°C after its production date. Keep out of direct sunlight and any heat source. Empty bottles must not be reused.

Approved for use in organic agriculture according to Reg. (EC) 889/2008: Annex I; (Micronutients, Microorganisms). Restriction: Micronutrient deficiency must be documented by testing or competent professional award.